illumina[®]

MiSeqDx[™] Cystic Fibrosis Clinical Sequencing Assay

The first FDA-cleared *in vitro* diagnostic next-generation sequencing assay to provide a complete view of the *CFTR* gene for increased confidence in cystic fibrosis variant detection.

Highlights

• Complete View of the CFTR Gene Capture all variants in the protein coding regions and intron/exon boundaries of the CFTR gene

Accurate Results

Deep coverage (> 3,000×) ensures detection accuracy with a Positive Agreement (PA) of 99.66%*

- Removed Demographic Bias Sequencing the *CFTR* gene removes the bias inherent in existing genotyping panels
- * PA is the number of samples with agreeing variant calls divided by the total number of samples with that variant as identified by the reference method.

Introduction

Cystic fibrosis (CF) affects approximately 70,000 children and adults worldwide¹. The disease appears when an individual inherits two mutated copies of the cystic fibrosis transmembrane receptor (*CFTR*) gene. Early diagnosis and treatment of CF can improve both survival and quality of life². Unfortunately, current molecular genotyping tests target a limited number of mutations, omitting variants that may have some clinical relevance. As a result, families may endure long periods of additional testing and patients may experience a delay in receiving needed treatment.

To overcome this challenge, Illumina offers the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay (Figure 1), the first FDA-cleared *in vitro* diagnostic (IVD) next-generation sequencing (NGS) test for obtaining a comprehensive view of the *CFTR* gene.



Figure 1: MiSeqDx Cystic Fibrosis Clinical Sequencing Assay. The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay is the first FDA-cleared NGS test to provide a comprehensive view of the *CFTR* gene.

Looking at the CFTR Sequence

Using Illumina NGS technology and sequencing by synthesis (SBS) chemistry, the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay sequences all protein coding regions and intron/exon boundaries of the *CFTR* gene. In addition, the assay detects two large deletions, two deep intronic mutations, and indels in homopolymeric regions such as the 2184delA deletion (Table 1). The assay also automatically detects polyTG/polyT variants. Looking at the full *CFTR* sequence minimizes the demographic bias inherent in existing CF genotyping panel tests. As a result, the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay can uncover rare mutations not accounted for in standard tests.

How it Works

The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay takes advantage of Illumina NGS technology and SBS chemistry to produce a reliable *CFTR* sequence. The streamlined sample preparation workflow replaces the multiple rounds of PCR required by Sanger sequencing with a single PCR step. For higher throughput, the assay enables multiplexing capabilities for simultaneous sequencing of 8 samples (Figure 2). Additionally, Illumina NGS is highly accurate and reproducible with \geq 99.7% overall agreement for all reported positions and a sample first pass rate of 99.7%.

Complete Kit Convenience

The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay includes everything needed for library preparation, sample multiplexing, and sequencing in a single kit. All reagents are packaged in a convenient ready-touse format, minimizing hands-on time and increasing uniformity in all tests.

Efficient Sample Preparation Increases Throughput

Library preparation begins with 250 ng genomic DNA (gDNA) isolated from a blood sample. The DNA is mixed with a pool of oligonucleotide probes. Each probe includes sequences designed to capture the designated variant and an adapter sequence used in a subsequent amplification reaction. The probes hybridize to the DNA, one upstream and one downstream of specific *CFTR* variants (Figure 3). A proprietary extension-ligation reaction extends across the region of interest, followed by ligation, to unite the two probes. This reaction creates a template strand and gives the assay excellent specificity.

To increase the number of samples analyzed in a single sequencing run, individual libraries are "tagged" with a unique identifier, or index. These unique sample-specific indices are added to each extensionligation template in a PCR amplification step. The final reaction product contains the *CFTR* variants with the necessary sequencing adapters and indices for sequencing on the MiSeqDx instrument. An automated three-read sequencing strategy identifies each tagged sample for individual downstream analysis. Using this approach, sample identification is highly accurate, maintaining the high integrity of the assay. Eight samples can be pooled in a single sequencing run.

Table 1: *CFTR* Regions Sequenced in the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay.

All protein coding regions

Intron/exon boundaries

Large deletions (exon 2,3 and exon 22,23)

Deep intronic mutations (1811+1.6Kb A>G, 3849+10Kb C>T)

Indels in homopolymeric regions

PolyTG/PolyT

PREPARE LIBRARIES

Generate sample libraries using the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay

Attach sample indices to libraries and amplify with one PCR step

Normalize and pool libraries

Hands-on time: 2 hr 10 min

SEQUENCE

Load pooled library on to MiSeqDx reagent cartridge

MiSeqDx instrument performs automated cluster generation and sequencing

Hands-on time: 5 min

ANALYZE

View report generated by the MiSeq Reporter software

Hands-on time: 5–30 min

Figure 2: Integrated Clinical Sequencing Assay Workflow. The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay offers a fully integrated end-to-end solution for sequencing the protein coding regions and intron/exon boundaries of the *CFTR* gene.




Figure 3: MiSeqDx Cystic Fibrosis Clinical Sequencing Assay Chemistry. The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay offers a highly specific method for sequencing the *CFTR* gene.

Widely Adopted NGS Platform

Illumina SBS technology is a widely adopted NGS platform in the sequencing community. Through massively parallel sequencing using a proprietary reversible terminator-based method, it enables detection of single bases as they are incorporated into growing DNA strands. A fluorescently labeled terminator is imaged as each dNTP (deoxyribonucleotide triphosphates: dATP, dCTP, dGTP, or dTTP) is added and then cleaved to allow incorporation of the next base. Since all four reversible terminator-bound dNTPs are present during each sequencing cycle, natural competition minimizes incorporation bias. The result is base-by-base sequencing that enables highly accurate data.

Easy Results Interpretation

Results from the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay are presented in an easy-to-read fashion that a board-certified molecular geneticist or equivalent can easily interpret. The assay report includes assay name, sample ID, dbSNP ID, HGVS nomenclature, and the call rates for each base (Figure 4). Call rates for each sample must be ≥ 99% to be considered valid.

Data Quality

The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay supports 8 samples per run, while providing excellent specificity and uniformity. To assess performance, accuracy, and reproducibility, studies were conducted using Sanger sequencing and PCR as reference methods. Total accuracy and reproducibility were ≥ 99.7%.

Summary

The MiSeqDx Cystic Fibrosis Clinical Sequencing Assay is the first FDA-cleared NGS diagnostic test for CF. Sequencing the *CFTR* gene provides a comprehensive genetic view not available using standard molecular genotyping panels. The additional data eliminates demographic bias and enables accurate detection of two large deletions, two deep intronic mutations, and indels in homopolymeric regions. This single assay can lead to faster identification of clinically relevant mutations.

	Reporter 2	2.30												
				1.	_									130329 M01018 0005 00000000-A1F
	and and and	alysis Info Sample Sh	ieet I	ogs Error	s									
	nples (48)											Search		Zoom 🕂 📶 🤇 117120009-117120240
	Sample ID	Sample Name	Clust		ster Align	Mismatch	No Call	Coverage	Het SNPs	Hom SNPs	Insertions	Deletions	Manifest	NA11497r 1250 1/CETR Ex1.chr7.117120007.117120211 tile 2.1
	NA11497_1250_1	NA11497_1250_1	2692		498/281094		0.00/0.08		1.0	1.1.25.6	0	2	CFTRManif *	3.5K
-	NA11497_1250_2	NA11497_1250_2	1821		233/178781		0.00/0.08		7	0	0	2	CFTRManif	3K- 2.5K
	NA11497_500_1	NA11497_500_1	2728		887/284254		0.00/0.08		7	0	0	12	CFTRManif	2.3K 2 2K-
	NA11497_500_2	NA11497_500_2	2685		581/260078		0.00/0.08		7	0	0	2	CFTRManif	≥ 1.5K
	NA11497_250_1	NA11497_250_1	1948		750/188309		0.00/0.08		7	0	0	2	CFTRManif	3 1K
	NA11497_250_2	NA11497_250_2	1388		142/134420		0.00/0.08		7	0	0	3	CFTRManif	500
	NA11497_100_1	NA11497_100_1	1479		927/141628	0.45/0.51	0.00/0.08		7	0	0	2	CFTRManif	40
	NA11497_100_2	NA11497_100_2			29/95894	0.46/0.52	0.00/0.08		r.			-	CFTRManif	35 MW monoun a monoun
	NA11497_50_1	NA11497_50_1	9942		13/94311	0.45/0.51	0.00/0.08		7	0	0	2	CFTRManif	30 V V V V V V V
	NA11497_50_2	NA11497_50_2	9205		30/87699	0.46/0.52	0.00/0.05		7	0	0	2	CFTRManif	5 20
	NA11497_25_1	NA11497_25_1	4743		77/45217	0.46/0.52	0.00/0.08	686.7	7	0	0	2	CFTRManif	Ö 15 10
	NA11497_25_2	NA11497_25_2	3857		08/34663	0.48/0.52	0.00/0.05		7	0	0	2	CFTRManif	5
	NA11497_10_1	NA11497_10_1	2382		83/22103	0.47/0.52	0.00/0.08		7	0	0	2	CFTRManif	0
	NA11497_10_2	NA11497_10_2	2172		65/19812	0.47/0.53	0.00/0.08		7	0	0	2	CFTRManif	3.5%
	NA11497_5_1	NA11497_5_1	1176	4 991	0/9819	0.48/0.52	0.00/0.05	149.5	7	0	0	2	CFTRManif 🗸	5 2.5K
•			-			_				_			•	Ę 1.5K
lar	jets (84)											Search		2 × 1K
Targe			Chr	Start Positio			uster PF	Mismatch	No Call		Hom SNPs	Insertions	Deletions	0 (0) (0) (0) (0)
	R_Ex1.chr7.11712000			117119879	117120		33/5884	2.45/0.32	0.00/0.04		0	0	0 ^	117120009 117120035 117120101 117120147 117120193 11712
	Ex1.chr7.11712000			117120009	117120		717/34345	0.53/0.82	0.00/0.05		1	0	0	Sample C G G C C C T A G C A G G G A C C C
CFTR_Intr1.chr7.117138297.117138437_tile_1.1			chr7	117138266	117138		476/75521	0.33/0.26	0.00/0.05	-	0	0	0	NA11407_1_1
CFTR_Ex2.chr7.117144282.117144434_tile_1.1			chr7 chr7	117144281	117144		1811/33437	0.32/0.35	0.00/0.05		0	0	0	NA11497_1_2 NA11497_1250_1
CFT	CFTR_Ex3.chr7.117149083.117149221_tile_1.1			117148938	117149		019/68070	0.37/0.37	0.00/0.05		1	0	1	NA11497_1250_1
CFTR_Ex3.chr7.117149083.117149221_tile_2.1			chr7	117149084	117149	287 52	999/52141	0.27/0.51	0.00/0.05	4	1	0	1	NA11497r_500_1
CFTR_Intr3.chr7.117159422.117159472_tile_1.1			chr7	117159379	117159	1553 10	7205/106663	0.45/0.30	0.00/0.05	2	0	0	0	N&11407+ Km 9
CFT	R_Ldel D1.chr7.11713	8297.117159472_tile_1.1	chr7	117138266	117138	473 0/	4	0.00/0.13	0.00/0.00	15	2	0	1	
CFT	R_Ex4.chr7.11717092	3.117171193_tile_1.1	chr7	117170799	117170	982 62	678/62494	0.49/0.29	0.00/0.05	3	0	0	0	Variants (1) Search
CFT	Ex4.chr7.11717092	3.117171193_tile_2.1	chr7	117170923	117171	108 57	416/56950	0.25/0.39	0.00/0.05	3	0	0	0	Sample ID Sample Name Chr Position Score Variant Type Call Frequency
CFT	Ex4.chr7.11717092	8.117171193_tile_3.1	chr7	117171055	117171	257 74	055/73895	0.38/0.27	0.00/0.08	3	1	0	0	NA11497r_1250_1 NA11497r_1250_1 chr7 117120141 3070 SNP G->[GiC] 0.49
CFTR_Ex5.chr7.117174250.117174446_tile_1.1			chr7	117174189	117174	390 39	794/39088	0.40/0.35	0.00/0.05	1	0	0	0	
CFT	Ex5.chr7.11717425	0.117174446_tile_2.1	chr7	117174337	117174	540 81	408/81027	0.30/0.48	0.00/0.05	4	0	0	0	
CFTR_Ex8.chr7.117175277.117175490_tile_1.1			chr7	117175148	117175	322 94	298/94112	0.37/0.38	0.00/0.04	1	1	0	0	
	F-8	7.117175490_tile_2.1	chr7	117175248	117175	442 75	276/72262	0.27/0.43	0.00/0.05	2	0	0	2 .	

Figure 4: Easy Visualization Using the MiSeq Reporter Software.

Learn More

To learn more about the MiSeqDx Cystic Fibrosis Clinical Sequencing Assay, visit www.illumina.com/cysticfibrosis.

References

- 1. Cystic Fibrosis Foundation (www.cff.org/AboutCF/Faqs/)
- 2. U.S. National Library of Medicine. PubMed Health (www.ncbi.nlm. nih.gov/pubmedhealth/PMH0001167/)

Ordering Information.

Product	Catalog No.
MiSeqDx Cystic Fibrosis Clinical Sequencing Assay (6 runs, up to 48 samples)	DX-102-1001

Intended Use

The Illumina MiSeqDx Cystic Fibrosis Clinical Sequencing Assay is a targeted sequencing *in vitro* diagnostic system that re-sequences the protein coding regions and intron/exon boundaries of the cystic fibrosis transmembrane conductance regulator (CFTR) gene in genomic DNA isolated from human peripheral whole blood specimens collected in K2EDTA. The test detects single nucleotide variants and small indels within the region sequenced, and additionally reports on two deep intronic mutations and two large deletions. The test is intended to be used on the Illumina MiSegDx instrument.

The test is intended to be used as an aid in the diagnosis of individuals with suspected cystic fibrosis (CF). This assay is most appropriate when the patient has an atypical or non-classic presentation of CF or when other mutation panels have failed to identify both causative mutations. The results of the test are intended to be interpreted by a board-certified clinical molecular geneticist or equivalent and should be used in conjunction with other available information including clinical symptoms, other diagnostic tests, and family history. This test is not indicated for use for stand-alone diagnostic purposes, fetal diagnostic testing, pre-implantation testing, carrier screening, newborn screening, or population screening.

Illumina • 1.800.809.4566 toll-free (U.S.) • +1.858.202.4566 tel • techsupport@illumina.com • www.illumina.com

FOR IN VITRO DIAGNOSTIC USE. CONTACT AN ILLUMINA REPRESENTATIVE FOR REGIONAL AVAILABILITY.

© 2015 Illumina, Inc. All rights reserved.

Illumina, MiSeqDx, and the pumpkin orange color are trademarks of Illumina, Inc. and/or its affiliate(s) in the U.S. and/or other countries. Pub. No. 670-2013-007 Current as of 26 August 2015 Document # 100000006361 v00

